

Hierarchy Problems in String Theory: The Power of Large Volume

Joseph P. Conlon (DAMTP, Cambridge)

Rutgers University, April 2007

This talk represents edited highlights of

[hep-th/0502058](#) (V. Balasubramanian, P. Berglund, JC, F. Quevedo)

[hep-th/0505076](#) (JC, F. Quevedo, K. Suruliz)

[hep-th/0602233](#) (JC)

[hep-th/0605141](#) (JC, F. Quevedo)

[hep-th/0609180](#) (JC, D.Cremades, F. Quevedo)

[hep-th/0610129](#) (JC, S. Abdussalam, F. Quevedo, K. Suruliz)

[hep-ph/0611144](#) (JC, D. Cremades)

[0704.xxxx \[hep-ph\]](#) (JC, C. Kom, K. Suruliz, B. Allanach, F. Quevedo)

Talk Structure

- Hierarchies in Nature
- String Phenomenology and Moduli Stabilisation
- Large Volume Models
- Axions
- Neutrino Masses
- Supersymmetric Soft Terms and Phenomenology
- Conclusions

Hierarchies in Nature

Nature likes hierarchies:

- The Planck scale, $M_P = 2.4 \times 10^{18} \text{GeV}$.
- The GUT/inflation scale, $M \sim 10^{16} \text{GeV}$.
- The axion scale, $10^9 \text{GeV} \lesssim f_a \lesssim 10^{12} \text{GeV}$
- The weak scale : $M_W \sim 100 \text{GeV}$
- The QCD scale $\Lambda_{QCD} \sim 200 \text{MeV}$ - understood!
- The neutrino mass scale, $0.05 \text{eV} \lesssim m_\nu \lesssim 0.3 \text{eV}$.
- The cosmological constant, $\Lambda \sim (10^{-3} \text{eV})^4$

These demand an explanation!

Hierarchies in Nature

In this talk I will enthusiastically advocate

- stabilised exponentially large extra dimensions ($\mathcal{V} \sim 10^{15} l_s^6$).
- an intermediate string scale $m_s \sim 10^{11} \text{GeV}$

as giving a natural, explanatory explanation for the axionic, weak and neutrino hierarchies.

The different hierarchies will come as different powers of the (large) volume.

String Phenomenology

- String theory lives in ten dimensions.
- The world is four-dimensional, so ten-dimensional string theory needs to be compactified.
- To preserve $\mathcal{N} = 1$ supersymmetry, we compactify on a Calabi-Yau manifold.
- The spectrum of light particles is determined by higher-dimensional topology.
- As part of the spectrum, string compactifications generically produce many uncharged scalar particles - **moduli** - that parametrise the size and shape of the extra dimensions.

Moduli Stabilisation

- Moduli are naively massless scalar fields which may take large classical vevs.
- They are uncharged and interact gravitationally.
- Such massless scalars generate unphysical fifth forces.
- The moduli need to be stabilised and given masses.
- Generating potentials for moduli is the field of **moduli stabilisation**.
- The large-volume models represent a particular (and appealing) moduli stabilisation scenario.

Moduli Stabilisation: Fluxes

- Flux compactifications involve non-vanishing flux fields on Calabi-Yau cycles.
- The fluxes carry a potential energy which depends on the geometry of the cycles.
- This energy generates a potential for the moduli associated with these cycles.
- In IIB compactifications, 3-form fluxes generate a superpotential

$$W = \int (F_3 + iSH_3) \wedge \Omega \equiv \int G_3 \wedge \Omega.$$

- This stabilises the dilaton and complex structure moduli.

Moduli Stabilisation: Fluxes

$$\hat{K} = -2 \ln (\mathcal{V}(T + \bar{T})) - \ln \left(i \int \Omega \wedge \bar{\Omega}(U) \right) - \ln (S + \bar{S}),$$

$$W = \int G_3 \wedge \Omega(S, U).$$

$$V = e^{\hat{K}} \left(\sum_{U,S} \hat{K}^{\alpha\bar{\beta}} D_{\alpha} W D_{\bar{\beta}} \bar{W} + \sum_T \hat{K}^{i\bar{j}} D_i W D_{\bar{j}} \bar{W} - 3|W|^2 \right)$$
$$= e^{\hat{K}} \left(\sum_{U,S} \hat{K}^{\alpha\bar{\beta}} D_{\alpha} W D_{\bar{\beta}} \bar{W} \right)$$

Stabilise S and U by solving $D_S W = D_U W = 0$.

Moduli Stabilisation: KKLT

$$\hat{K} = -2 \ln(\mathcal{V}) - \ln \left(i \int \Omega \wedge \bar{\Omega} \right) - \ln(S + \bar{S}),$$
$$W = \int G_3 \wedge \Omega + \sum_i A_i e^{-a_i T_i}.$$

Non-perturbative effects (D3-instantons / gaugino condensation) allow the T -moduli to be stabilised by solving $D_T W = 0$.

For consistency, this requires

$$W_0 = \left\langle \int G_3 \wedge \Omega \right\rangle \ll 1.$$

Moduli Stabilisation: Large-Volume

$$\hat{K} = -2 \ln \left(\mathcal{V} + \frac{\xi}{g_s^{3/2}} \right) - \ln \left(i \int \Omega \wedge \bar{\Omega} \right) - \ln (S + \bar{S}),$$

$$W = \int G_3 \wedge \Omega + \sum_i A_i e^{-a_i T_i}.$$

We include the leading α' corrections to the Kähler potential.

This leads to dramatic changes in the large-volume vacuum structure.

Moduli Stabilisation: Large-Volume

The simplest model $\mathbb{P}^4_{[1,1,1,6,9]}$ has two Kähler moduli.

$$\mathcal{V} = \left(\left(\frac{T_b + \bar{T}_b}{2} \right)^{3/2} - \left(\frac{T_s + \bar{T}_s}{2} \right)^{3/2} \right) \equiv \left(\tau_b^{3/2} - \tau_s^{3/2} \right).$$

If we compute the scalar potential, we get for $\mathcal{V} \gg 1$,

$$V = \frac{\sqrt{\tau_s} a_s^2 |A_s|^2 e^{-2a_s \tau_s}}{\mathcal{V}} - \frac{a_s |A_s W_0| \tau_s e^{-a_s \tau_s}}{\mathcal{V}^2} + \frac{\xi |W_0|^2}{g_s^{3/2} \mathcal{V}^3}.$$

The minimum of this potential can be found analytically.

Moduli Stabilisation: Large-Volume

Integrate out τ_s :

$$V = \underbrace{\frac{\sqrt{\tau_s} a_s^2 |A_s|^2 e^{-2a_s \tau_s}}{\mathcal{V}} - \frac{a_s |A_s W_0| \tau_s e^{-a_s \tau_s}}{\mathcal{V}^2}}_{\text{}} + \frac{\xi |W_0|^2}{g_s^{3/2} \mathcal{V}^3}.$$

$$V = - \frac{|W_0|^2 (\ln \mathcal{V})^{3/2}}{\mathcal{V}^3} + \frac{\xi |W_0|^2}{g_s^{3/2} \mathcal{V}^3}.$$

A minimum exists at

$$\mathcal{V} \sim |W_0| e^{c/g_s}, \quad \tau_s \sim \ln \mathcal{V}.$$

This minimum is **non-supersymmetric AdS** and at **exponentially large volume**.


Moduli Stabilisation: Large-Volume

- The large volume $\mathcal{V} \gg 1$ controls the α' expansion and justifies the inclusion only of the leading term.
- The correction used arises from the 10d \mathcal{R}^4 term.
- This is dominant - other ten-dimensional higher-derivative terms are suppressed by higher powers of \mathcal{V} , e.g.

$$\int d^{10}x G_3^4 \mathcal{R}^2 \sim \mathcal{V}^{-4/3} \int d^{10}x \mathcal{R}^4.$$

- Quantum loop corrections are also suppressed compared to the \mathcal{R}^4 term.

Moduli Stabilisation: Large-Volume



Moduli Stabilisation: Large-Volume

The mass scales present are:

- Planck scale: $M_P = 2.4 \times 10^{18} \text{ GeV}$.
- String scale: $M_S = \frac{M_P}{\sqrt{\mathcal{V}}} \sim 10^{11} \text{ GeV}$.
- KK scale $M_{KK} = \frac{M_P}{\mathcal{V}^{2/3}} \sim 10^9 \text{ GeV}$.
- Gravitino mass $m_{3/2} = \frac{M_P}{\mathcal{V}} \sim 30 \text{ TeV}$.
- Small Kähler moduli
 $m_{\tau_s} \sim m_{3/2} \ln(M_P/m_{3/2}) \sim 1000 \text{ TeV}$.
- Complex structure moduli $m_U \sim m_{3/2} \sim 30 \text{ TeV}$.
- Soft terms $m_{susy} \sim \frac{m_{3/2}}{\ln(M_P/m_{3/2})} \sim 1 \text{ TeV}$.
- Volume modulus $m_{\tau_b} \sim \frac{M_P}{\mathcal{V}^{3/2}} \sim 1 \text{ MeV}$.

Summary of Large Volume Models

- The stabilised volume is naturally exponentially large.
- The large volume lowers the string scale and gravitino mass through

$$m_s = \frac{M_P}{\sqrt{\mathcal{V}}}, \quad m_{3/2} = \frac{M_P}{\mathcal{V}}.$$

- With $m_s = 10^{11}$ GeV, the weak hierarchy can be naturally generated through TeV-scale supersymmetry.
- The minimum is non-supersymmetric and generates gravity-mediated soft terms - **more later....**

Axions

- Axions are a well-motivated solution to the strong CP problem.
- The QCD Lagrangian is

$$\mathcal{L}_{QCD} = \frac{1}{g^2} \int d^4x F_{\mu\nu}^a F^{a,\mu\nu} + \theta \int F^a \wedge F^a.$$

- The strong CP problem is that naively $\theta \in (-\pi, \pi)$ while experimentally, $|\theta| \lesssim 10^{-10}$.
- The axionic (Peccei-Quinn) solution is to promote θ to a dynamical field, $\theta(x)$.

Axions

- The canonical Lagrangian for θ is


$$\mathcal{L} = \frac{1}{2} \partial_\mu \theta \partial^\mu \theta + \int \frac{\theta}{f_a} F^a \wedge F^a.$$

- f_a is called the axionic decay constant.
- Constraints on supernova cooling and direct searches imply $f_a \gtrsim 10^9 \text{ GeV}$.
- Avoiding the overproduction of axion dark matter prefers $f_a \lesssim 10^{12} \text{ GeV}$.
- There exists an axion ‘allowed window’,

$$10^9 \text{ GeV} \lesssim f_a \lesssim 10^{12} \text{ GeV}.$$

Axions

- For D7 branes, the axionic coupling comes from the RR form in the brane Chern-Simons action.
- The axion decay constant f_a measures the coupling of the axion to matter.



Axions

- The coupling of the axion to matter is a local coupling and does not see the overall volume.
- This coupling can only see the string scale, and so

$$f_a \sim m_s \sim \frac{M_P}{\sqrt{\mathcal{V}}}.$$

(This is confirmed by a full analysis)

- This generates the axion scale,

$$f_a \sim \frac{M_P}{\sqrt{\mathcal{V}}} \sim 10^{11} \text{ GeV}.$$

Neutrino Masses

- The theoretical origin of neutrino masses is a mystery. Experimentally

$$0.05\text{eV} \lesssim m_\nu^H \lesssim 0.3\text{eV}.$$

- This corresponds to a Majorana mass scale

$$M_{\nu_R} \sim 3 \times 10^{14}\text{GeV}.$$

- Equivalently, this is the suppression scale Λ of the dimension five Standard Model operator

$$\mathcal{O} = \frac{1}{\Lambda} H H L L.$$

Neutrino Masses

- In the supergravity MSSM, consider the superpotential operator

$$\frac{\lambda}{M_P} H_2 H_2 L L \in W,$$

where λ is dimensionless.

- This gives rise to the Lagrangian terms,

$$\tilde{K}_{H_2} \partial_\mu H_2 \partial^\mu H_2 + \tilde{K}_L \partial_\mu L \partial^\mu L + e^{\hat{K}/2} \frac{\lambda}{M_P} H_2 H_2 L L.$$

- This corresponds to the *physical* coupling

$$e^{\hat{K}/2} \frac{\lambda}{M_P} \frac{\langle H_2 H_2 \rangle L L}{(\tilde{K}_{H_2} \tilde{K}_{H_2} \tilde{K}_L \tilde{K}_L)^{\frac{1}{2}}}.$$

Neutrino Masses

- We have the coupling

$$e^{\hat{K}/2} \frac{\lambda}{M_P} \frac{\langle H_2 H_2 \rangle LL}{(\tilde{K}_{H_2} \tilde{K}_{H_2} \tilde{K}_L \tilde{K}_L)^{\frac{1}{2}}}.$$

- Once the Higgs receives a vev, this generates neutrino masses.
- However, to compute the mass scale we need to know \hat{K}_{H_2} and \hat{K}_L ...

Computing the Kähler Metric

- I now describe new techniques for computing the matter metric $\tilde{K}_{\alpha\bar{\beta}}$ for chiral matter fields on a Calabi-Yau.
- These techniques will apply to chiral bifundamental D7/D7 matter in IIB compactifications.
- I will compute the modular weights of the matter metrics from the modular dependence of the (physical) Yukawa couplings.

I start by reviewing Yukawa couplings in supergravity.

Yukawa Couplings in Supergravity

- In supergravity, the Yukawa couplings arise from the Lagrangian terms (Wess & Bagger)

$$\begin{aligned}\mathcal{L}_{kin} + \mathcal{L}_{yukawa} &= \tilde{K}_\alpha \partial_\mu C^\alpha \partial^\mu \bar{C}^{\bar{\alpha}} + e^{\hat{K}/2} \partial_\beta \partial_\gamma W \psi^\beta \psi^\gamma \\ &= \tilde{K}_\alpha \partial_\mu C^\alpha \partial^\mu \bar{C}^{\bar{\alpha}} + e^{\hat{K}/2} Y_{\alpha\beta\gamma} C^\alpha \psi^\beta \psi^\gamma\end{aligned}$$

(This assumes diagonal matter metrics, but we can relax this)

- The matter fields are not canonically normalised.
- The *physical* Yukawa couplings are given by

$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma)^{\frac{1}{2}}}.$$

Computing $\tilde{K}_{\alpha\bar{\beta}}$ (I)

$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma)^{\frac{1}{2}}}.$$

- We know the modular dependence of \hat{K} :

$$\hat{K} = -2 \ln(\mathcal{V}) - \ln \left(i \int \Omega \wedge \bar{\Omega} \right) - \ln(S + \bar{S})$$

- We compute the modular dependence of \tilde{K}_α from the modular dependence of $\hat{Y}_{\alpha\beta\gamma}$. We work in a power series expansion and determine the leading power λ ,

$$\tilde{K}_\alpha \sim (T + \bar{T})^\lambda k_\alpha(\phi) + (T + \bar{T})^{\lambda-1} k'_\alpha(\phi) + \dots$$

Computing $\tilde{K}_{\alpha\bar{\beta}}$ (II)

$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_{\alpha}\tilde{K}_{\beta}\tilde{K}_{\gamma})^{\frac{1}{2}}}.$$

- We are unable to compute $Y_{\alpha\beta\gamma}$. If $Y_{\alpha\beta\gamma}$ depends on a modulus T , knowledge of $\hat{Y}_{\alpha\beta\gamma}$ gives no information about the dependence $\tilde{K}_{\alpha\bar{\beta}}(T)$.
- Our results will be restricted to those moduli that do not appear in the superpotential.
- This holds for the T -moduli (Kähler moduli) in IIB compactifications. In perturbation theory,

$$\partial_{T_i} Y_{\alpha\beta\gamma} \equiv 0.$$

Computing $\tilde{K}_{\alpha\bar{\beta}}$ (III)


$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma)^{\frac{1}{2}}}.$$

- We know $\hat{K}(T)$.
- If we can compute $\hat{Y}_{\alpha\beta\gamma}(T)$, we can then deduce $\tilde{K}_\alpha(T)$.
- We can compute $\hat{Y}_{\alpha\beta\gamma}(T)$ for IIB compactifications through wavefunction overlap.

We now describe the computation of $\hat{Y}_{\alpha\beta\gamma}$ for bifundamental matter on a stack of magnetised D7-branes.

The brane geometry

- We consider a stack of (magnetised) branes all wrapping an identical cycle.
- Chiral fermions stretch between differently magnetised branes.



Computing the physical Yukawas $\hat{Y}_{\alpha\beta\gamma}$

- We use a simple computational technique:
- Physical Yukawa couplings are determined by the triple overlap of normalised wavefunctions.
- These wavefunctions can be computed (in principle) by dimensional reduction of the brane action. At low energies, this DBI action reduces to Super Yang-Mills.

Computing the physical Yukawas $\hat{Y}_{\alpha\beta\gamma}$

Consider the higher-dimensional term

$$\mathcal{L} \sim \int_{\mathbb{M}_4 \times \Sigma} d^n x \sqrt{g} \bar{\lambda} \Gamma^i (A_i + \partial_i) \lambda$$

Dimensional reduction of this gives the kinetic terms

$$\bar{\psi} \partial \psi$$

and the Yukawa couplings

$$\bar{\psi} \phi \psi$$

- The physical Yukawa couplings are set by the combination of the above!

Kinetic Terms

$$\mathcal{L} \sim \int_{\mathbb{M}_4 \times \Sigma} d^n x \sqrt{g} \bar{\lambda} \Gamma^i (A_i + \partial_i) \lambda$$

- Dimensional reduction gives

$$\lambda = \psi_4 \otimes \psi_6, \quad A_i = \phi_4 \otimes \phi_6.$$

and the reduced kinetic terms are

$$\left(\int_{\Sigma} \psi_6^\dagger \psi_6 \right) \int_{\mathbb{M}_4} \bar{\psi}_4 \Gamma^\mu (A_\mu + \partial_\mu) \psi_4$$

- Canonically normalised kinetic terms require

$$\int_{\Sigma} \psi_6^\dagger \psi_6 = 1.$$

Yukawa Couplings

$$\mathcal{L} \sim \int_{\mathbb{M}_4 \times \Sigma} d^n x \bar{\lambda} \Gamma^i (A_i + \partial_i) \lambda$$

- Dimensional reduction gives the four-dimensional interaction

$$\left(\int_{\Sigma} \bar{\psi}_6 \Gamma^I \phi_{I,6} \psi_6 \right) \int_{\mathbb{M}_4} d^4 x \phi_4 \bar{\psi}_4 \psi_4.$$

- The physical Yukawa couplings are determined by the (normalised) overlap integral

$$\hat{Y}_{\alpha\beta\gamma} = \left(\int_{\Sigma} \bar{\psi}_6 \Gamma^I \phi_{I,6} \psi_6 \right).$$

The Result

$$\int_{\Sigma} \psi_6^\dagger \psi_6 = 1, \quad \hat{Y}_{\alpha\beta\gamma} = \left(\int_{\Sigma} \bar{\psi}_6 \Gamma^I \phi_{I,6} \psi_6 \right).$$

- For canonical kinetic terms,

$$\psi_6(y) \sim \frac{1}{\sqrt{\text{Vol}(\Sigma)}}, \quad \Gamma^I \phi_{I,6} \sim \frac{1}{\sqrt{\text{Vol}(\Sigma)}},$$

and so

$$\hat{Y}_{\alpha\beta\gamma} \sim \text{Vol}(\Sigma) \cdot \frac{1}{\sqrt{\text{Vol}(\Sigma)}} \cdot \frac{1}{\sqrt{\text{Vol}(\Sigma)}} \cdot \frac{1}{\sqrt{\text{Vol}(\Sigma)}} \sim \frac{1}{\sqrt{\text{Vol}(\Sigma)}}$$

- This gives the scaling of $\hat{Y}_{\alpha\beta\gamma}(T)$.

Application: Large-Volume Models

- The physical Yukawa coupling

$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma)^{1/2}}$$

is local and thus independent of \mathcal{V} .

- As $\hat{K} = -2 \ln \mathcal{V}$, we can deduce *simply from locality* that

$$\tilde{K}_\alpha \sim \frac{1}{\mathcal{V}^{2/3}}.$$

- As this is for a Calabi-Yau background, this is already non-trivial!

Application: Large-Volume Models

- We can go further. With all branes wrapping the same 4-cycle,

$$\hat{Y}_{\alpha\beta\gamma} = e^{\hat{K}/2} \frac{Y_{\alpha\beta\gamma}}{(\tilde{K}_\alpha \tilde{K}_\beta \tilde{K}_\gamma)^{1/2}} \sim \frac{1}{\sqrt{\tau_s}}.$$

- We can then deduce that

$$\tilde{K}_{\alpha\bar{\beta}} \sim \frac{\tau_s^{1/3}}{\mathcal{V}^{2/3}} k_{\alpha\bar{\beta}}(U, \bar{U}).$$

- We also have the dependence on τ_s !

Neutrino Masses

- We have the coupling

$$e^{\hat{K}/2} \frac{\lambda}{M_P} \frac{\langle H_2 H_2 \rangle LL}{(\tilde{K}_{H_2} \tilde{K}_{H_2} \tilde{K}_L \tilde{K}_L)^{\frac{1}{2}}}.$$

- Once the Higgs receives a vev, this generates neutrino masses.
- However, to compute the mass scale we need to know \tilde{K}_{H_2} and \tilde{K}_L ...done!



$$\tilde{K}_{\alpha\bar{\beta}} \sim \frac{\tau_s^{1/3}}{\mathcal{V}^{2/3}} k_{\alpha\bar{\beta}}(\phi).$$

Neutrino Masses

- Using the large-volume result $\tilde{K}_\alpha \sim \frac{\tau_s^{1/3}}{\mathcal{V}^{2/3}}$, this becomes

$$\frac{\lambda \mathcal{V}^{1/3}}{\tau_s^{2/3} M_P} \langle H_2 H_2 \rangle LL.$$

- Use $\mathcal{V} \sim 10^{15}$ (to get $m_{3/2} \sim 1\text{TeV}$) and $\tau_s \sim 10$:

$$\frac{\lambda}{10^{14} \text{GeV}} \langle H_2 H_2 \rangle LL$$

- With $\langle H_2 \rangle = \frac{v}{\sqrt{2}} = 174\text{GeV}$, this gives

$$m_\nu = \lambda(0.3 \text{ eV}).$$

SUSY Breaking and Soft Terms

The MSSM is specified by its matter content (that of the supersymmetric Standard Model) plus soft breaking terms. These are

- Soft scalar masses, $m_i^2 \phi_i^2$
- Gaugino masses, $M_a \lambda^a \lambda^a$,
- Trilinear scalar A-terms, $A_{\alpha\beta\gamma} \phi^\alpha \phi^\beta \phi^\gamma$
- B-terms, $BH_1 H_2$.

The soft terms break supersymmetry explicitly, but do not reintroduce quadratic divergences into the Lagrangian.

We want to compute these soft terms for the large volume models.

SUSY Breaking and Soft Terms

- For gravity mediation, the computation of soft terms starts by expanding the supergravity K and W in terms of the matter fields C^α ,

$$W = \hat{W}(\Phi) + \mu(\Phi)H_1H_2 + \frac{1}{6}Y_{\alpha\beta\gamma}(\Phi)C^\alpha C^\beta C^\gamma + \dots,$$

$$K = \hat{K}(\Phi, \bar{\Phi}) + \tilde{K}_{\alpha\bar{\beta}}(\Phi, \bar{\Phi})C^\alpha C^{\bar{\beta}} + [ZH_1H_2 + h.c.] + \dots,$$

$$f_a = f_a(\Phi).$$

- Given this expansion, the computation of the physical soft terms is straightforward.
- The function $\tilde{K}_{\alpha\bar{\beta}}(\Phi)$ is crucial in computing soft terms, as it determines the normalisation of the matter fields.

SUSY Breaking and Soft Terms

- Soft scalar masses m_{ij}^2 and trilinears $A_{\alpha\beta\gamma}$ are given by

$$\begin{aligned} \tilde{m}_{\alpha\bar{\beta}}^2 &= (m_{3/2}^2 + V_0) \tilde{K}_{\alpha\bar{\beta}} \\ &\quad - \bar{F}^{\bar{m}} F^n \left(\partial_{\bar{m}} \partial_n \tilde{K}_{\alpha\bar{\beta}} - (\partial_{\bar{m}} \tilde{K}_{\alpha\bar{\gamma}}) \tilde{K}^{\bar{\gamma}\delta} (\partial_n \tilde{K}_{\delta\bar{\beta}}) \right) \\ A'_{\alpha\beta\gamma} &= e^{\hat{K}/2} F^m \left[\hat{K}_m Y_{\alpha\beta\gamma} + \partial_m Y_{\alpha\beta\gamma} \right. \\ &\quad \left. - \left((\partial_m \tilde{K}_{\alpha\bar{\rho}}) \tilde{K}^{\bar{\rho}\delta} Y_{\delta\beta\gamma} + (\alpha \leftrightarrow \beta) + (\alpha \leftrightarrow \gamma) \right) \right]. \end{aligned}$$

- *Any* physical prediction for the soft terms requires a knowledge of $\tilde{K}_{\alpha\bar{\beta}}$ for chiral matter fields.
- Fortunately, we just described how to compute $\tilde{K}_{\alpha\bar{\beta}}$.

SUSY Breaking and Soft Terms

- We can compute the soft terms for the large-volume models...

(details skipped)

... in the dilute-flux approximation we get

$$\begin{aligned}M_i &= \frac{F^s}{2\tau_s} \equiv M, \\m_{\alpha\bar{\beta}} &= \frac{M}{\sqrt{3}} \tilde{K}_{\alpha\bar{\beta}}, \\A_{\alpha\beta\gamma} &= -M \hat{Y}_{\alpha\beta\gamma}, \\B &= -\frac{4M}{3}.\end{aligned}$$

Soft Terms: Flavour Universality

- These soft terms are flavour-universal.
- Gravity mediation ‘generically’ gives non-universal soft terms: both flavour and susy breaking are Planck-scale physics, and so the susy-breaking sector will be sensitive to flavour.
- However -

Soft Terms: Flavour Universality

- These soft terms are flavour-universal.
- Gravity mediation ‘generically’ gives non-universal soft terms: both flavour and susy breaking are Planck-scale physics, and so the susy-breaking sector will be sensitive to flavour.
- However - this is an effective field theory argument.
- In string theory, we have Kähler (T) and complex structure (U) moduli. These are **decoupled** at leading order.

$$\mathcal{K} = -2 \ln(\mathcal{V}(T)) - \ln \left(i \int \Omega \wedge \bar{\Omega}(U) \right) - \ln(S + \bar{S}).$$

- The kinetic terms for T and U fields do not mix.

Soft Terms: Flavour Universality

- Due to the shift symmetry $T \rightarrow T + i\epsilon$, the T moduli make no perturbative appearance in the superpotential.
- It is the U moduli that source flavour...

$$W = \dots + \frac{1}{6} Y_{\alpha\beta\gamma}(U) C^\alpha C^\beta C^\gamma + \dots$$

- ...and the T moduli that break supersymmetry,

$$D_T W \neq 0, F^T \neq 0, \quad D_U W = 0, F^U = 0.$$

- At leading order, susy breaking (Kähler moduli) and flavour (complex structure moduli) decouple.
- The soft terms are automatically flavour-universal.

Soft Terms: Spectra


- Magnetic fluxes on the brane world-volumes are necessary for chirality.
- They also alter e.g. the gauge kinetic functions

$$f_a = \frac{T}{4\pi} \rightarrow f_a = \frac{T}{4\pi} + h_a(F)S.$$

- This affects the soft terms; we study the effect on the resulting sparticle spectrum by generating soft terms perturbed about those for the dilute flux case.
- We generate many such spectra, with high-scale soft terms allowed to fluctuate by $\pm 20\%$.

Soft Terms: Spectra

- We run the soft terms to low energy using SoftSUSY:



Soft Terms: Spectra

- The spectrum is more compressed than a typical mSUGRA spectra: the squarks tend to be lighter and the sleptons heavier.
- This arises because the RG running starts at the intermediate rather than GUT scale.
- The gaugino mass ratios are

$$M_1 : M_2 : M_3 = 1.5 \rightarrow 2 : 2 : 6.$$

This can be distinguished from both mSUGRA and mirage mediation.

A new scale....

- Large-volume models robustly predict the existence of a new gravitationally coupled scalar particle χ with mass $\sim 1\text{MeV}$.
- This particle can decay via $\chi \rightarrow 2\gamma$ and $\chi \rightarrow e^+e^-$.
- One can show

$$Br(\chi \rightarrow e^+e^-) \sim \frac{\ln(M_P/m_{3/2})^2}{20} Br(\chi \rightarrow 2\gamma)$$

- The e^+e^- decay is strongly preferred - may be of interest with regard to the 511 keV line....

Large Volumes are Power-ful

In large-volume models, an exponentially large volume appears naturally ($\mathcal{V} \sim e^{\frac{c}{g_s}}$). The scales that naturally appear are

- Susy-breaking: $m_{soft} \sim \frac{M_P}{\mathcal{V}} \sim 10^3 \text{ GeV}$
- Axions: $f_a \sim \frac{M_P}{\sqrt{\mathcal{V}}} \sim 10^{11} \text{ GeV}$
- Neutrinos/dim-5 operators: $\Lambda \sim \frac{M_P}{\mathcal{V}^{1/3}} \sim 10^{14} \text{ GeV}$
- All three scales (plus flavour universality) are yoked in an attractive fashion.
- The origin of all three hierarchies is the exponentially large volume.

