The Coming Era of New Physics at the Large Hadron Collider.

Talk to the Rutgers Society of Physics Students March 30, 2011

Amitabh Lath




So what's a collider do, exactly?

 Colliders (accelerators) have been essential tools to understand the structure of nature.




- In a remarkably short period of time, our understanding of the basic building blocks of nature has changed.
 - Several times!
- The technique used is called scattering.







Scattering Basics



Scattering Basics



What you can see depends on *resolution*




Large wavelength, low energy, target looks like a big blob.

Short wavelength, Higher resolution




Small wavelength, high energy, constituents of targets become visible


THE ELECTROMAGNETIC SPECTRUM



Scattering at higher energies (shorter wavelengths)

- Microscope to Short Distances
- New Substructure and Forces





What are things made of ~1900


Dmitrii Mendeleev:

Periodic Table of Elements.

 Everything is made up of mixtures of pure elements (ATOMS).

JJ Thompson: electron.


 ATOMS can eject these small, negatively charged bits that are also the carrier of electric current.



- 58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	HARMAN LAND	100000000000000000000000000000000000000				Lr

Picture of the Atom ~1900

Thompson plum pudding model of the atom




Copyright @ 2000 Benjamin/Cummings, an imprint of Addison Wesley Longman, Inc.

Picture of the Atom ~1900

Thompson plum pudding model of the atom


Today, the plum pudding model is synonymous with the flat earth theory, but in its time it answered some important questions.

How can a heavy, overall neutral object emit light charged particles when heated (ionization?)




Copyright © 2000 Benjamin/Cummings, an imprint of Addison Wesley Longman, Inc.

Rutherford Destroys the Plum Pudding Model




Rutherford Destroys the Plum Pudding Model






Rutherford



Marsden


Brilliant example of undergraduate contribution to particle physics. Tradition continues to this day!

What Rutherford Expected




Projectiles (very fast He nuclei called *alpha particles*) will be slightly deflected by gold atoms

What Rutherford Saw



Occasionally (rarely) the projectile scattered at huge angles. Beginning of 20th century physics.

What Rutherford Saw



Occasionally (rarely) the projectile scattered at huge angles. Beginning of 20th century physics.


Things started small...

The first cyclotron...



...and got bigger


Lawrence next to the Berkeley cyclotron ...discovery of pions. (~simultaneously with cosmic rays)



...and even bigger

Princeton-Stanford accelerator (Prin-Stan).

500 MeV, 2-ring electron-electron machine




Not all were circular...

The Stanford Linear Accelerator.


Delivered 15 GeV electrons on target.



End Station A




What Kendall, Friedman, and Taylor expected...



NO large angle scattering.

What Kendall, Friedman, and Taylor actually saw

Electrons from the Linear accelerator




Proton (target)

Large angle scattering can happen.

And did! Quarks!

What Kendall, Friedman, and Taylor actually saw



Proton (target)

Quarks!

The 1990 Nobel Prize




OBSERVED BEHAVIOR OF HIGHLY INELASTIC ELECTRON-PROTON SCATTERING

M. Breidenbach, J. I. Friedman, and H. W. Kendall

Department of Physics and Laboratory for Nuclear Science,*
Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

and

E. D. Bloom, D. H. Coward, H. DeStaebler, J. Drees, L. W. Mo, and R. E. Taylor Stanford Linear Accelerator Center,† Stanford, California 94305 (Received 22 August 1969)



Seen (Bjorken scaling, aka scattering off quarks)

y toevio

FIG. 1. $(d^2\sigma/d\Omega dE')/\sigma_{\rm Mott}$, in GeV⁻¹, vs q^2 for W=2, 3, and 3.5 GeV. The lines drawn through the data are meant to guide the eye. Also shown is the cross section for elastic e-p scattering divided by $\sigma_{\rm Mott}$, $(d\sigma/d\Omega)/\sigma_{\rm Mott}$, calculated for θ =10°, using the dipole form factor. The relatively slow variation with q^2 of the inelastic cross section compared with the elastic cross section is clearly shown.


PRL Vol 23, No 16, (1969)

Aside: Antimatter




Dirac predicted antimatter as a consequence of combining relativity and quantum mechanics.

Anderson found it (an antielectron, called positron) in cosmic ray data.



Even bigger Colliders


(now with matter-antimatter collisions!)



Most were just "let's build it and see what's there". Most discoveries in the field were indeed surprises!


SppS was built to see W/Z for which there were precise predictions

LHC is built to explore electroweak energy scale which it completely covers



So what is everything made of?


Status as of 2011




Everything in the universe is made of the 1st generation particles.

Quarks bind together with the strong force to make familiar particles such as protons and neutrons.

So why isn't this good enough?





antimatter

Go back 100 years...

$$\nabla \cdot \mathbf{E} = \rho/\epsilon_0$$

$$\nabla \cdot \mathbf{B} = 0$$


$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

$$\nabla \times \mathbf{B} = \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} + \mu_0 \mathbf{j}_c$$

Isn't this good enough?

Even before QED, we knew that classical electrodynamics could not be the whole story . . .


The classical theory predicts its own demise with an infinite electron self-energy



(This is a recurring and important theme)

Nonsensical predictions, and solutions


Fermi theory of the 1930's



This process violates unitarity at high energies

What do we do?


Modify the diagram to cancel the divergence



the W boson

(observed at CERN in 1983)


Nonsensical predictions, and solutions cont.



But now this process violates unitarity at high energies!

What do we do?

Introduce another diagram that cancels the divergence



the Z boson

(also observed at CERN in 1983)

Nonsensical predictions, and solutions cont.

 W^{+} W^{-} W^{-} W^{+} W^{-} W^{+} W^{-} W^{+} W^{-} W^{+} W^{-} W^{+} W^{+}

But now these processes violate unitarity at high energies!

What do we do?


Introduce *other* diagrams to cancel the divergence

 $W^{+} \sim \sim W^{+}$ $W^{-} \sim W^{+}$ $W^{+} \sim W^{+}$ $W^{-} \sim W^{+}$

the Higgs boson

Nonsensical predictions, and solutions cont.

Thus far we have <u>no direct evidence</u> for the Higgs boson*



If the Higgs exists, these types of processes violate unitarity at high energies unless a parameter is "unnaturally" fine-tuned ("fine-tuning problem")

What do we do?

Introduce other ____ diagrams to cancel the divergence without fine-tuning

supersymmetry
strong dynamics
extra dimensions

Symmetries

Symmetries are the Central Organizing Principle in Our Understanding of Fundamental Physics !!!

Spatial Translation Invariance (Momentum)

Time Translation Invariance (Energy)


Spatial Rotation Invariance (Angular

Momentum)


Space-Time Rotation / Boost Invariance



(Lorentz, ..., Einstein)



Supersymmetry




Every SM particle gets a *super*-partner to cancel inconvenient divergences. Superpartners of fermions are bosons (add an "s": selectron, squark...) Superpartners of bosons are fermions (add an "ino": wino, gluino, photino...)

Dark matter candidates!

Superpartners more massive than SM, but should be around ~TeV or so...

Hunting grounds of the LHC!


The Large Hadron Collider



Large Hadron Collider



Radio Frequency Klystron Cavities 1232 Super Conducting Dipole Magnets



T = 1.9 K I = 12000 Amps


B = 8.3 Tesla E = 7 MJ / Dipole

½ nanogram in Beam -

Kinetic Energy of 100,000 Ton
Aircraft Carrier at Cruising Speed


Vue d'ensemble des expériences LHC. 31211212 LHC - B Point 8 **CERN** ATLAS Point 1 **ALICE** Point 2 CMS Point 5 SPS ATLAS LHC - B LEP/LHC **CMS** Photothèque - E540 - V10/09/97


High Energy Collider Detectors



CMS Detector Transverse Slice


Particle Identification





LHC and CMS operations in 2010.


About 47pb⁻¹ delivered by LHC and ~43pb⁻¹ of data collected by CMS. Overall data taking efficiency ~92%. 6pb⁻¹ of data integrated in a good fill. Excellent performance in coping with more than 5 order of magnitude increase in instantaneous luminosity.




Average fraction of operational channels per CMS sub-system still ~99%. A few problems here and there. Last few days of pp running tested 50ns filling scheme. Vacuum (e-cloud) worse than at 150ns. 75ns vacuum much better. 800 bunches OK.

LHC and CMS operations in 2010.

About 47pb⁻¹ delivered by LHC and ~43pb⁻¹ of data collected by CMS. Overall data taking efficiency ~92%. 6pb⁻¹ of data integrated in a good fill. Excellent performance in coping with more than 5 order of magnitude increase in instantaneous luminosity.



Average frac post operational channels per CMS sub-system still ~99%. A few problems here and there. Last few days of pp running tested 50ns filling scheme. Vacuum (e-cloud) worse than at 150ns. 75ns vacuum much better. 800 bunches OK.



Standard Model Versus New Physics Processes

Yesterday's Discovery
Today's Background
Tomorrow's Calibration

100,000,000 Top Quarks / yr Design Luminosity


Opportunity for Precision Top Physics ...


Huge Background to New Physics Searches

May Have to Enhance Recorded S/B by 10⁽⁴⁻⁵⁾
Skill of the Physics Analysis

Typically Look in Low Background Channels

– But what if nature doesn't live there?






Beautiful ZZ event seen in CMS data. First found by RU postdoc R. Gray.

 $\mu_0 + \mu_1$: 92.15 GeV (total(Z) p_T 26.5 GeV, ϕ -3.03), $\mu_2 + \mu_3$: 92.24 GeV (total(Z) p_T 29.4 GeV, ϕ +.06), $\mu_0 + \mu_2$: 70.12 GeV (total p_T 27 GeV), $\mu_3 + \mu_1$: 83.1 GeV (total p_T 26.1 GeV).


Invariant Mass of 4µ: 201 GeV

Multilepton SUSY Search




Many other models for new physics have multiple leptons as decay signatures.


This one analysis also puts bounds on **Slepton Co-NLSP** (where the sleptons are close in mass to the lightest SUSY Particle) models with **leptonic R-Parity volations.** Both of these give 4 leptons In the final state.



Diphoton + missing momentum



Jet Physics



Summary

- The energy frontier has always been surprising.
 - No one expected the nucleus, heavy leptons, mesons, partons, charm...
- Incredibly beautiful data coming out of LHC detectors.
 - Almost makes you forget how long it took...
 - Granularity of CMS/ATLAS much finer than CDF/D0 → entirely new physics techniques.
- We have entered the era of new physics. The 2011-2012 run is being called the "discovery run".
 - Is there a higgs? Will we create Dark Matter?
 - Whatever happens, it probably will not be what we expect.
 - Will SUSY etc go the way of S-Matrix theory?
- To the students in the audience: JOIN US!