


1 Consider the unit sphere  $S^2$ . Embedded in three dimensional Euclidean space  $\mathbb{R}^3$  with cartesian coordinates, this is the set  $\{(x, y, z) | x^2 + y^2 + z^2 = 1\}$ , but as it is a two dimensional space, we usually describe it with spherical polar coordinates  $\theta, \phi$ , fixing the radius  $r = 1$ , with  $z = \cos \theta$  and  $(x, y) = \sin \theta(\cos \phi, \sin \phi)$ . The coordinates  $\theta$  and  $\phi$  are well-defined, however, only for  $\theta \neq 0$  and  $\neq \pi$ . So we have a chart  $C$  for  $\mathcal{U}$  the unit sphere with the north and south pole removed.

The unit sphere has a natural metric induced from the Euclidean metric of  $\mathbb{R}^3$ , with which we are very familiar:

$$(ds)^2 = (d\theta)^2 + \sin^2 \theta (d\phi)^2, \quad \text{or} \quad g_{\theta\theta} = 1, g_{\phi\phi} = \sin^2 \theta, g_{\theta\phi} = 0.$$

- (a) Calculate the Christoffel symbol  $\Gamma^j_{kl}$
- (b) The shortest path (on the sphere) between two points is determined by the geodesic equations. Find the equations which determine  $\theta(s)$  and  $\phi(s)$ , where the  $s$  parameter is the path length. Show that one of these second order equations is essentially the conservation of  $L_z$ .
- (c) It is easier to guess the solution from the knowledge that the answer is a great circle, the intersection of the sphere with a plane through the origin, rather than solving the equations from scratch. Show that this is correct, satisfying the geodesic equations.

2 Consider the unit sphere  $S^2$  as the set of point  $\{(x, y, z) | x^2 + y^2 + z^2 = 1\}$  in  $\mathbb{R}^3$ . This is really a two dimensional manifold, so it needs to be considered as such. Consider the two open sets  $\mathcal{U}_N$  which is the sphere minus the South pole, and  $\mathcal{U}_S$ , the sphere minus the North pole. Define the charts  $C_N$  and  $C_S$



by considering the equatorial plane  $\mathcal{E}$ . For each point  $\mathcal{P} \in \mathcal{U}_N$  on the sphere, draw straight lines (in  $\mathbb{R}^3$ ) through the South pole and  $\mathcal{P}$ , and define  $\phi_N(\mathcal{P})$  to be the point  $(x_N, y_N)$  at which the line intersects the equatorial plane. Similarly define  $\phi_S(\mathcal{P})$  to be the point in equatorial plane on the straight line through the

- (a) Find the coordinates of  $\phi_N(\mathcal{P})$  in terms of the  $(x, y, z)$  of  $\mathcal{P}$  in  $\mathbb{R}^3$ , and the inverse relation.
- (b) Do the same for  $\phi_S(\mathcal{P})$ , and find the transition function

$$\phi_{SN} = \phi_S \circ \phi_N^{-1} = \phi_S(\phi_N^{-1}(x_N, y_N)).$$

- (c) The metric on the sphere is  $(ds)^2 = (d\theta)^2 + \sin^2 \theta (d\phi)^2$ . Find the metric on the chart  $C_N$ . What is the relation to the ordinary Euclidean metric on the plane?

World North Pole Stereographic Projection Map

