0.1 Wronskian

A set of n vectors \vec{V}_j is linearly independent if the only set $a_j \in F$ such that $\sum_j a_j \vec{V}_j = 0$ is $a_j = 0$ for all j. If the vectors are in an n dimensional space and they are given in terms of a set of basis vectors, we have $\sum_j a_j (V_j)_k = 0$ for all k, which means the matrix $M_{kj} = (V_j)_k$ annihilates the vector a_j , and the \vec{V}_j are linearly independent if and only if det M = 0.

If we have a set of n linearly independent solutions $y_k(x)$ to an n'th order ordinary linear differential equation, the Wronskian is defined as the determinant of the matrix of j'th derivatives, $j = 0, \ldots, n-1$ of the n functions y,

$$W(x) := \det \frac{d^{\ell-1}y_k}{dx^{\ell-1}} = \sum_{i_1,\dots,i_n} \epsilon_{i_1,\dots,i_n} \prod_{j=0}^{n-1} \frac{d^j y_{i_j}}{dx^j},$$

where by d^0y/dx^0 we just mean y. If we had a set $a_k(x)$ such that

$$\sum_{k} a_k(x) \frac{d^{\ell-1}y_k}{dx^{\ell-1}}(x) = 0 \text{ for all } \ell, \text{ without the } a_k(x) \text{ all vanishing, we would}$$

have a linear dependence among our n solutions, because the n'th derivative would also vanish, as each y_k satisfies the equation. So linear independence tells us the determinant does not vanish.

If we differentiate W, we have

$$\frac{dW}{dx} = \sum_{i_1, \dots, i_n} \epsilon_{i_1, \dots, i_n} \sum_{\ell=0}^{n-1} \prod_{j=0}^{n-1} \frac{d^{j+\delta_{j\ell}} y_{i_j}}{dx^{j+\delta_{j\ell}}}.$$

Note that unless $\ell = n - 1$, the terms in the product with $j = \ell$ and the terms with $j = \ell + 1$ are now identical, except for interchanging the indices on y, so the ϵ kills them, and we have only the contribution from $\ell = n - 1$, which is

$$\frac{dW}{dx} = \sum_{i_1,\dots,i_n} \epsilon_{i_1,\dots,i_n} \frac{d^n y_{i_n}}{dx^n} \prod_{j=0}^{n-2} \frac{d^j y_{i_j}}{dx^j}.$$

Now if our differential equation is

$$\frac{d^n y}{dx^n}(x) + P(x)\frac{d^{n-1} y}{dx^{n-1}}(x) + \sum_{\ell=0}^{n-2} Q_{\ell}(x)\frac{d^{\ell} y}{dx^{\ell}}(x) = 0$$

we can substitute the values of $d^n y_{i_n}/dx^n$,

$$\frac{dW}{dx} = -\sum_{i_1,\dots,i_n} \epsilon_{i_1,\dots,i_n} \prod_{j=0}^{n-2} \frac{d^j y_{i_j}}{dx^j} \times \left(P(x) \frac{d^{n-1} y_{i_n}}{dx^{n-1}}(x) + \sum_{\ell=1}^{n-2} Q_{\ell}(x) \frac{d^{\ell} y_{i_n}}{dx^{\ell}}(x) \right) \\
= -\sum_{i_1,\dots,i_n} \epsilon_{i_1,\dots,i_n} \prod_{j=0}^{n-1} \frac{d^j y_{i_j}}{dx^j} P(x) - \sum_{i_1,\dots,i_n} \sum_{\ell=1}^{n-2} Q_{\ell}(x) \epsilon_{i_1,\dots,i_n} \frac{d^{\ell} y_{i_n}}{dx^{\ell}} \prod_{j=0}^{n-2} \frac{d^j y_{i_j}}{dx^j}$$

Note that in the terms involving Q_{ℓ} , the term before the product matches one of the terms in the product, except for interchange $i_n \leftrightarrow i_{\ell}$, and so they are killed by the ϵ . Also notice the term multiplying -P(x) is just W(x), so we have the first order differential equation

$$\frac{dW}{dx} = -P(x)W(x).$$