

# Comment on Integrals over Grassman Variables

Joel A. Shapiro

With the rule for integrating over a Grassman variable,

$$\int d\theta f(\theta) = B, \quad \text{where } f(\theta) = A + \theta B,$$

we can define nested integrals, so that if  $f$  is a function of  $n$   $\theta$ 's, we may define

$$\int d^n \theta f(\{\theta_i\}) = \int d\theta_n d\theta_{n-1} \dots d\theta_1 f(\{\theta_i\}) = \text{coefficient of } \theta_1 \theta_2 \dots \theta_n.$$

Note the order of the  $d\theta$ 's in  $d^n \theta$ .

Note that an equivalent rule for how  $\int d\theta$  acts on  $f(\theta)$  is to define it as: *take the  $\theta$  derivative!* Thus it would appear that

$$\int d\theta_n d\theta_{n-1} \dots d\theta_1 = \frac{\partial}{\partial \theta_n} \frac{\partial}{\partial \theta_{n-1}} \dots \frac{\partial}{\partial \theta_1},$$

and in fact people often think about the integral as really being a derivative. Some caution is needed, however.

We will assume for the moment that our  $\theta$ 's are real. As we are allowed to multiply them by real numbers and add them, as a vector space, we might consider a change in variables,

$$\theta'_i = A_{ij} \theta_j,$$

where  $A$  is a real matrix. If we reexpress  $f$  in terms of those variables,

$$f(\{\theta_i\}) = f'(\{\theta'_i\}),$$

the coefficients  $c_n$  for  $f$  and  $c'_n$  for  $f'$  are related by

$$c'_n (A_{1,j_1} \theta_{j_1}) (A_{2,j_2} \theta_{j_2}) \dots (A_{n,j_n} \theta_{j_n}) = c_n \theta_1 \theta_2 \dots \theta_n.$$

Rearranging the  $\theta$ 's on the left gives  $\epsilon_{j_1, j_2, \dots, j_n}$ . so

$$c_n = c'_n \epsilon_{j_1, j_2, \dots, j_n} \prod_{i=1}^n A_{i, j_i} = c'_n \det A.$$

If we assume the infinitesimal  $d$  (the exterior derivative) is linear over the ordinary numbers, so  $d\theta'_i = A_{ij} d\theta_j$ , then  $d^n \theta' = \det A d^n \theta$  exactly as for the product of  $n$   $\theta$ 's, and we find

$$\int d^n \theta' f'(\{\theta'\}) = \det A \int d^n \theta f(\{\theta\}) = c_n \det A = c'_n (\det A)^2,$$

which is not what we would have found by assuming the  $\theta$ 's were as valid as the  $\theta$ 's, unless  $(\det A) = \pm 1$ , as would be the case for an orthogonal matrix. Of course restricting to orthogonal transformations is reasonable.

But to define complex  $\theta$ 's Peskin and Schroeder define

$$\theta = \frac{\theta_1 + i\theta_2}{\sqrt{2}}, \quad \theta^* = \frac{\theta_1 - i\theta_2}{\sqrt{2}}, \quad (1)$$

which is of the form above with a complex unitary matrix  $A$  which is not orthogonal and has determinant  $-i$ . Thus the rest of the steps leading to their 9.66 are dubious.

If we suppose linearity of  $d$ , (1) implies

$$d\theta = \frac{d\theta_1 + id\theta_2}{\sqrt{2}}, \quad d\theta^* = \frac{d\theta_1 - id\theta_2}{\sqrt{2}}, \quad (2)$$

but then

$$\int d\theta \theta = \int \frac{d\theta_1 + id\theta_2}{\sqrt{2}} \frac{\theta_1 + i\theta_2}{\sqrt{2}} = \frac{1}{2} \int d\theta_1 \theta_1 - d\theta_2 \theta_2 = 0,$$

while

$$\int d\theta \theta^* = \int \frac{d\theta_1 + id\theta_2}{\sqrt{2}} \frac{\theta_1 - i\theta_2}{\sqrt{2}} = \frac{1}{2} \int d\theta_1 \theta_1 + d\theta_2 \theta_2 = 1.$$

and also  $\int d\theta^* \theta = 1$ . That seems somewhat strange but might be suitable ( $\theta$  and  $\theta^*$  are complex so as fields would be considered charged, with opposite charge, while the integral has lost any Grassman charge and should be neutral). But it would also make

$$\int d\theta^* d\theta \theta \theta^* = - \int d\theta^* d\theta \theta^* \theta = - \int d\theta^* \left( \int d\theta \theta^* \right) \theta = - \int d\theta^* \theta = -1,$$

in disagreement with P&S.