

Lecture 7: Dirac and Weyl Fields Sept. 26, 2013

Copyright©2005 by Joel A. Shapiro

We have seen that we expect to construct our field theory from fields which transform “simply” under Poincaré transformations, with

$$U(\Lambda)\phi_a(x)U^{-1}(\Lambda) = D_{ab}(\Lambda^{-1})\phi_b(\Lambda x), \quad (1)$$

where D is a finite dimensional representation of the Lorentz group. We also saw that such representations are in fact products of representations of two $SO(3)$ groups generated by $\vec{L}_\pm = \frac{1}{2}(\vec{J} \pm i\vec{K})$. Thus in general there are two spins, s_\pm and the field has two indices, the eigenvalues of $\vec{L}_{\pm z}$ respectively. The derivative terms of (L6 Eq. 10) can be simplified

$$\begin{aligned} \vec{\theta} \cdot \vec{J}^\mu_\nu x^\nu \partial_\mu &= +\frac{1}{2}\epsilon_{ijk}\theta_i (\mathcal{L}_{jk})^\mu_\nu x^\nu \partial_\mu = i\vec{\theta} \cdot (\vec{x} \times \vec{\nabla}), \\ \vec{\kappa} \cdot \vec{K}^\mu_\nu x^\nu \partial_\mu &= \kappa_i (\mathcal{L}_{0j})^\mu_\nu x^\nu \partial_\mu = -i\vec{\kappa} \cdot \vec{x} \partial_0 - it\vec{\kappa} \cdot \vec{\nabla}. \end{aligned}$$

Then the operators \vec{J} and \vec{K} have commutators with fields given by

$$\begin{aligned} [\vec{\theta} \cdot \vec{J}, \phi_{m_+, m_-}(x)] &= -D_{m_+, m'_+}^A(\vec{\theta} \cdot \vec{L})\phi_{m'_+, m_-}(x) \\ &\quad - D_{m_-, m'_-}^B(\vec{\theta} \cdot \vec{L})\phi_{m_+, m'_-}(x) \\ &\quad + i\vec{\theta} \cdot (\vec{x} \times \vec{\nabla})\phi_{m_+, m_-}(x) \\ [\vec{\kappa} \cdot \vec{K}, \phi_{m_+, m_-}(x)] &= +iD_{m_+, m'_+}^A(\vec{\kappa} \cdot \vec{L})\phi_{m'_+, m_-}(x) \\ &\quad - iD_{m_-, m'_-}^B(\vec{\kappa} \cdot \vec{L})\phi_{m_+, m'_-}(x) \\ &\quad + -i\vec{\kappa} \cdot \vec{x} \dot{\phi}_{m_+, m_-}(x) - it\vec{\kappa} \cdot \vec{\nabla}\phi_{m_+, m_-}(x). \end{aligned}$$

In particular, we considered a field (whose name I will now change to ψ_R , which transforms with $A = \frac{1}{2}, B = 0$, and we saw that

$$[\vec{J}, \psi_{Rm}(x)] = -\frac{1}{2}\vec{\sigma}_{mm'}\psi_{Rm'}(x) + i\vec{x} \times \vec{\nabla}\psi_{Rm}(x).$$

We will consider this field further, but before we do, let us also note that the Poincaré algebra $[\mathbf{L}_{\alpha\beta}, \mathbf{P}_\nu] = -ig_{\alpha\nu}\mathbf{P}_\beta + ig_{\beta\nu}\mathbf{P}_\alpha$ means

$$[\mathbf{J}_i, \mathbf{P}_j] = \frac{1}{2}\epsilon_{iab}[\mathbf{L}_{ab}, \mathbf{P}_j] = i\epsilon_{ijk}\mathbf{P}_k, \quad [\mathbf{J}_i, \mathbf{P}_0] = 0. \quad (2)$$

$$[\mathbf{K}_i, \mathbf{P}_j] = [\mathbf{L}_{0i}, \mathbf{P}_j] = -i\delta_{ij}\mathbf{P}_0, \quad [\mathbf{K}_i, \mathbf{P}_0] = [\mathbf{L}_{0i}, \mathbf{P}_0] = -i\mathbf{P}_i. \quad (3)$$

The $(\frac{1}{2}, 0)$ Field ψ_R

We are going to look for scalar combinations of fields, in order to construct a Lagrangian density \mathcal{L} . The coordinate derivative terms will work out as they should for any representation, so in what follows I am going to drop the derivative terms, with just a warning (+d.t.) that I have done so.

First suppose that ψ_R transforms with $A = \frac{1}{2}, B = 0$, so

$$[\mathbf{J}_i, \psi_R] = -\frac{1}{2}\sigma_i\psi_R, \quad [\mathbf{K}_i, \psi_R] = +i\frac{1}{2}\sigma_i\psi_R \quad (+d.t.).$$

Hermitean conjugate gives, as \mathbf{J} and \mathbf{K} are hermitean operators on the hilbert space, but the 2×2 representation of \mathbf{K} is not,

$$[\mathbf{J}_i, \psi_R^\dagger] = \frac{1}{2}\psi_R^\dagger\sigma_i, \quad [\mathbf{K}_i, \psi_R^\dagger] = +i\frac{1}{2}\psi_R^\dagger\sigma_i, \quad (+d.t.)$$

What can we make that is quadratic in ψ and its hermitian conjugate, and how do these terms transform?

$$[\mathbf{J}_i, \psi_R^\dagger\psi_R] = \frac{1}{2}\psi_R^\dagger\sigma_i\psi_R - \frac{1}{2}\psi_R^\dagger\sigma_i\psi_R = 0 \quad (+d.t.)$$

$$[\mathbf{J}_i, \psi_R^\dagger\sigma_j\psi_R] = \frac{1}{2}\psi_R^\dagger[\sigma_i, \sigma_j]\psi_R = i\epsilon_{ijk}\psi_R^\dagger\sigma_k\psi_R \quad (+d.t.)$$

$$[\mathbf{K}_i, \psi_R^\dagger\psi_R] = i\psi_R^\dagger\sigma_i\psi_R \quad (+d.t.)$$

$$[\mathbf{K}_i, \psi_R^\dagger\sigma_j\psi_R] = \frac{i}{2}\psi_R^\dagger\{\sigma_i, \sigma_j\}\psi_R = i\delta_{ij}\psi_R^\dagger\psi_R \quad (+d.t.)$$

Combining with (2) we see that $\psi_R^\dagger\psi_R$, $\psi_R^\dagger\sigma_j\psi_R$ and $\psi_R^\dagger P_0\psi_R$ commute with \mathbf{J}_i . We seek a combination which commutes with \mathbf{K}_i as well.

$$\begin{aligned} [K_i, \psi_R^\dagger\psi_R P_0] &= [K_i, \psi_R^\dagger\psi_R] P_0 + \psi_R^\dagger\psi_R [K_i, P_0] \\ &= i\psi_R^\dagger\sigma_i\psi_R P_0 - i\psi_R^\dagger\psi_R P_i \end{aligned} \quad (4)$$

$$\begin{aligned} \left[K_i, \sum_j \psi_R^\dagger\sigma_j\psi_R P_j \right] &= \left[K_i, \sum_j \psi_R^\dagger\sigma_j\psi_R \right] P_j + i \sum_j \psi_R^\dagger\sigma_j\psi_R [K_i, P_j] \\ &= i\psi_R^\dagger\psi_R P_i - i\psi_R^\dagger\sigma_i\psi_R P_0 \end{aligned} \quad (5)$$

so

$$\left[\mathbf{K}_i, \psi_R^\dagger\psi_R P_0 + \sum_j \psi_R^\dagger\sigma_j\psi_R P_j \right] = 0 \quad (+d.t.).$$

The $(0, \frac{1}{2})$ Field ψ_L

On the other hand, suppose ψ_L transforms with $A = 0, B = \frac{1}{2}$, so

$$D(J_i - iK_i) = \frac{1}{2}\sigma_i, \quad D(J_i + iK_i) = 0 \quad \implies \quad D(J_i) = \frac{1}{2}\sigma_i, \quad D(K_i) = +i\frac{1}{2}\sigma_i,$$

The commutations with \mathbf{J}_i are therefore all the same, while the one of the fields with \mathbf{K}_i are reversed, but not those of \mathbf{K} with P . So now the first terms in the final expressions in (4) and (5) have their signs reversed, and the combination which is a scalar is

$$\psi_L^\dagger \psi_L P_0 - \sum_i \psi_L^\dagger \sigma_i \psi_L P_i.$$

Notice there is no invariant we can make from just ψ_R without a momentum, or from just ψ_L without a momentum, but if we mix ψ_R with ψ_L , we see $\psi_R^\dagger \psi_L$ commutes with \mathbf{J} as before, and also

$$[\mathbf{K}_i, \psi_R^\dagger \psi_L] = i\frac{1}{2}\psi_R^\dagger \sigma_i \psi_L - i\frac{1}{2}\psi_L^\dagger \sigma_i \psi_R = 0,$$

so $\psi_R^\dagger \psi_L$ is an invariant. Similarly $\psi_L^\dagger \psi_R$ is invariant..

1 Invariant Lagrangians

The momentum transforms the same way a derivative does, so we see that the Hermitean quadratic invariants we can form from ψ_R and ψ_L are

$$\begin{aligned} & i\psi_R^\dagger \partial_0 \psi_R + i\psi_R^\dagger \vec{\sigma} \cdot \vec{\nabla} \psi_R \\ & i\psi_L^\dagger \partial_0 \psi_L - i\psi_L^\dagger \vec{\sigma} \cdot \vec{\nabla} \psi_L \\ & \psi_R^\dagger \psi_L + \psi_L^\dagger \psi_R \\ \text{and } & i\psi_R^\dagger \psi_L - i\psi_L^\dagger \psi_R \end{aligned}$$

The only one which involves only ψ_R is the first, and if we vary with respect to ψ_R^\dagger , we get the equation of motion

$$i\partial_0 \psi_R + i\vec{\sigma} \cdot \vec{\nabla} \psi_R = 0.$$

Multiplying by $-i(\partial_0 - \vec{\sigma} \cdot \vec{\nabla})$ gives¹

$$\begin{aligned} 0 &= (\partial_0 - \vec{\sigma} \cdot \vec{\nabla})(\partial_0 + \vec{\sigma} \cdot \vec{\nabla})\psi_R = (\partial_0^2 - \sigma_i \partial_i \sigma_j \partial_j)\psi_R \\ &= (\partial_0^2 - \frac{1}{2}\{\sigma_i, \sigma_j\} \partial_i \partial_j)\psi_R = (\partial_0^2 - \delta_{ij} \partial_i \partial_j)\psi_R \\ &= (\partial_0^2 - \vec{\nabla}^2)\psi_R = \partial^\mu \partial_\mu \psi_R. \end{aligned}$$

In the second line we have used the fact that $\partial_i \partial_j = \partial_j \partial_i$ to replace $\sigma_i \sigma_j$ by half the anticommutator, which we then evaluate to a Kronecker delta. We see the result is that ψ_R obeys the Klein-Gordon equation, but with zero mass. The same is true for the second lagrangian, with only ψ_L . Only by including a term with a mixture of ψ_R and ψ_L can we create a mass.

Let's define²

$$\sigma_R^\mu = (1, \sigma_i), \quad \text{and} \quad \sigma_L^\mu = (1, -\sigma_i).$$

Then we can write the first two lagrangian densities as $i\psi_R^\dagger \sigma_R^\mu \partial_\mu \psi_R$ and $i\psi_L^\dagger \sigma_L^\mu \partial_\mu \psi_L$, and the equations of motion from them individually as $\sigma_R^\mu \partial_\mu \psi_R = 0$ and $\sigma_L^\mu \partial_\mu \psi_L = 0$.

If, however, we take a combination to form the lagrangian,

$$\mathcal{L} = i\psi_R^\dagger \sigma_R^\mu \partial_\mu \psi_R + i\psi_L^\dagger \sigma_L^\mu \partial_\mu \psi_L - m(\psi_R^\dagger \psi_L + \psi_L^\dagger \psi_R),$$

we get the equations of motion

$$\begin{aligned} i\sigma_R^\mu \partial_\mu \psi_R - m\psi_L &= 0 \\ i\sigma_L^\mu \partial_\mu \psi_L - m\psi_R &= 0 \end{aligned} \quad \text{or} \quad \begin{pmatrix} -m & i\sigma_R^\mu \partial_\mu \\ i\sigma_L^\mu \partial_\mu & -m \end{pmatrix} \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} = 0.$$

Because we are mostly interested in massive fields, we will prefer to consider ψ_L and ψ_R as parts of a four component field. Define

$$\psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} \quad \text{and} \quad \gamma^\mu = \begin{pmatrix} 0 & \sigma_R^\mu \\ \sigma_L^\mu & 0 \end{pmatrix}$$

¹Properties of $\vec{\sigma}$: $\sigma_j = \sigma_j^\dagger$; $\sigma_i \sigma_j = \delta_{ij} + i\epsilon_{ijk} \sigma_k$, so $\{\sigma_i, \sigma_j\} = 2\delta_{ij}$ and $[\sigma_i, \sigma_j] = 2i\epsilon_{ijk} \sigma_k$. The usual representation, which we will assume, is

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

²The book, and indeed everyone else who defines these, uses σ^μ for what I call σ_R^μ and $\bar{\sigma}^\mu$ for what I call σ_L^μ . But that notation is not ideal.

which means

$$\gamma^0 = \begin{pmatrix} 0 & \mathbb{1}_{2 \times 2} \\ \mathbb{1}_{2 \times 2} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix}.$$

Then the equation above becomes

$$(i\gamma^\mu \partial_\mu - m)\psi = 0.$$

This is known as the Dirac equation.

A good part of learning how to calculate scattering amplitudes for fermions is becoming agile with the algebra of the γ matrices. From

$$\gamma^\mu \gamma^\nu = \begin{pmatrix} \sigma_R^\mu \sigma_L^\nu & 0 \\ 0 & \sigma_L^\mu \sigma_R^\nu \end{pmatrix}$$

we see that

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu} \times \mathbb{1}_{4 \times 4}, \quad (6)$$

which of course means $\gamma^{02} = 1$, $\gamma^{i2} = -1$.

Premultiplying the equation of motion $0 = (i\gamma^\mu \partial_\mu - m)\psi$ by $-(i\gamma^\nu \partial_\nu + m)$ we see that

$$\begin{aligned} 0 &= -(i\gamma^\nu \partial_\nu + m)(i\gamma^\mu \partial_\mu - m)\psi = \left(\frac{1}{2} \{\gamma^\nu, \gamma^\mu\} \partial_\nu \partial_\mu + m^2\right) \psi \\ &= (g^{\nu\mu} \partial_\nu \partial_\mu + m^2) \psi = (\partial^\mu \partial_\mu + m^2) \psi, \end{aligned}$$

so the Dirac equation implies the Klein-Gordon equation with mass m , but has additional information in it.

The γ matrices will prove to be much more often used than our σ_L^μ and σ_R^μ , so we need to reexpress our Lagrangian in terms of them. Notice that $\gamma^0 \gamma^\mu = \begin{pmatrix} \sigma_L & 0 \\ 0 & \sigma_R \end{pmatrix}$ so our lagrangian can be written

$$\mathcal{L} = i\psi^\dagger \gamma^0 \gamma^\mu \partial_\mu \psi - m\psi^\dagger \gamma^0 \psi.$$

That looks very strange, not even covariant, but the reason for this is that ψ^\dagger does not transform as we might expect, because ψ transforms under a representation $D(\Lambda) = \Lambda_{\frac{1}{2}}$ (or $(\frac{1}{2}, 0) + (0, \frac{1}{2})$), which is **not** a unitary representation of the Lorentz group, because \vec{L}_\pm involves $i\vec{K}$. Under $\psi \rightarrow \Lambda_{\frac{1}{2}}\psi$,

we have $\psi^\dagger \psi \rightarrow \psi^\dagger \Lambda_{\frac{1}{2}}^\dagger \Lambda_{\frac{1}{2}} \psi$, and if $\Lambda_{\frac{1}{2}}$ were unitary we would have $\Lambda_{\frac{1}{2}}^\dagger = \Lambda_{\frac{1}{2}}^{-1}$, and $\psi^\dagger \psi$ would be invariant, as it appears. But this is not the case.

What is $\Lambda_{\frac{1}{2}}$? It turns out there is a simple expression for its generators in terms of

$$S^{\mu\nu} = \frac{i}{4} [\gamma^\mu, \gamma^\nu].$$

From the anticommutation relations of the gammas (6) simple algebraic manipulations show that $D(L^{\mu\nu}) = S^{\mu\nu}$ obeys the Lorentz algebra commutation relations, and thus is a representation. In fact, from

$$S^{ij} = \frac{1}{2} \epsilon_{ijk} \begin{pmatrix} \sigma_k & 0 \\ 0 & \sigma_k \end{pmatrix}, \quad S^{0j} = -\frac{i}{2} \begin{pmatrix} \sigma_j & 0 \\ 0 & -\sigma_j \end{pmatrix},$$

we see that this is exactly how ψ transforms, or rather that

$$\Lambda_{\frac{1}{2}} \left(e^{-\frac{i}{2} \omega^{\mu\nu} \mathbf{L}_{\mu\nu}} \right) = e^{-\frac{i}{2} \omega^{\mu\nu} S_{\mu\nu}}.$$

Now notice that $\gamma_0^{-1} \gamma^\mu \gamma_0 = \gamma_\mu = (\gamma^\mu)^\dagger$, which means that $\gamma_0^{-1} S_{\mu\nu}^\dagger \gamma_0 = S_{\mu\nu}$ and $\gamma_0^{-1} \Lambda_{\frac{1}{2}}^\dagger \gamma_0 = \Lambda_{\frac{1}{2}}^{-1}$. Thus if we define $\bar{\psi} := \psi^\dagger \gamma_0$, under a Lorentz transformation

$$\bar{\psi} \rightarrow \psi^\dagger \Lambda_{\frac{1}{2}}^\dagger \gamma_0 = \psi^\dagger \gamma_0 \gamma_0^{-1} \Lambda_{\frac{1}{2}}^\dagger \gamma_0 = \bar{\psi} \Lambda_{\frac{1}{2}}^{-1},$$

so $\bar{\psi}\psi$ is invariant, and so is $\bar{\psi}\gamma^\mu\partial_\mu\psi$. Thus we can rewrite the free Dirac lagrangian density as

$$\mathcal{L}_{\text{Dirac}} = \bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi.$$