

13th ~~Feb~~ Feb, 2014

Course: Biophysics
Problem Set: 2

Due: 20th Feb, 2014

3.1 White-collar crime

- You are a city inspector. You go undercover to a bakery and buy 30 loaves of bread marked 500 g. Back at the lab you weigh them and find their masses to be 493, 503, 486, 489, 501, 498, 507, 504, 493, 487, 495, 498, 494, 490, 494, 490, 497, 503, 498, 495, 503, 496, 492, 492, 495, 498, 490, 490, 497, and 482 g. You go back to the bakery and issue a warning. Why?
- Later you return to the bakery (this time, they know you). They sell you 30 more loaves of bread. You take them home, weigh them, and find their masses to be 504, 503, 503, 503, 501, 500, 500, 501, 505, 501, 501, 500, 508, 503, 503, 500, 503, 501, 500, 502, 502, 501, 503, 501, 501, 502, 503, 501, 502, and 500 g. You're satisfied, because all the loaves weigh at least 500 g. But your boss reads your report and tells you to go back and close the shop down. What did she notice that you missed?

3.4 Photons

Section 3.3.3 reviewed Muller's and Timoféeff's empirical results that the rate of induced mutations is proportional to the radiation exposure. Not only X-rays can induce mutations; even ultraviolet light will work (that's why you wear sunblock). To get a feeling for what is so shocking about these results, notice that they imply that there's no "safe," or threshold, dose level. The amount of damage (probability of damaging a gene) is directly proportional to the total radiation exposure. Extrapolating to the smallest possible dose, we must conclude that even a single photon of UV light has the ability to cause permanent genetic damage to a skin cell and its progeny. (Photons are the packets of light mentioned in Section 1.5.3.)

- Somebody tells you that a single ultraviolet photon carries an energy equivalent of about 10 electron volts (eV, see Appendix B). You propose a damage mechanism: A photon delivers that energy into a volume the size of the cell nucleus and heats it up; then the increased thermal motion knocks the chromosomes apart in some

way. Is this a reasonable proposal? Why or why not? [Hint: Use Equation 1.2, and the definition of calorie found just below it, to calculate the temperature change.]

- Turning the result around, suppose that that photon's energy is delivered to a small volume L^3 and heats it up. We might suspect that if it heats up the region to boiling, this change could disrupt any genetic message contained in that volume. How small must L be for this amount of energy to heat that volume up to boiling (from 30°C to 100°C)? What could we conclude about the size of a gene if this proposal were correct?