

Physics 464/511 Homework #2

Due: Sept. 26, 2016 at 5:00 P. M.

1 [5 pts] Evaluate

$$\frac{1}{3} \int_S \vec{r} \cdot d\vec{\sigma}$$

over the unit cube defined by the point $(0,0,0)$ and the unit intercepts on the x -, y -, and z -axes. Note that

- (a) $\vec{r} \cdot d\vec{\sigma}$ is zero for three of the surfaces and
- (b) each of the three remaining surfaces contributes the same amount to the integral.

2 [5 pts] Show that

$$\frac{1}{3} \int_S \vec{r} \cdot d\vec{\sigma} = V$$

where V is the volume enclosed by the closed surface S .

3 [5 pts] An *algebra* consists of a vector space A over a field F , together with a binary operation of multiplication on the set A of vectors, ($A \times A \rightarrow A$) such that for all $a \in F$ and $\alpha, \beta, \gamma \in A$, the following are satisfied:

- (A) $(a\alpha)\beta = a(\alpha\beta) = \alpha(a\beta)$
- (B) $(\alpha + \beta)\gamma = \alpha\gamma + \beta\gamma$
- (C) $\alpha(\beta + \gamma) = \alpha\beta + \alpha\gamma$

If, in addition, (D): $\alpha(\beta\gamma) = (\alpha\beta)\gamma$, A is an associative algebra.

- (a) Show that linear transformations on a finite-dimensional vector space V into itself (endomorphisms) form an associative algebra A .
- (b) Define the commutator of two elements T_1 and T_2 of A by $[T_1, T_2] := T_1T_2 - T_2T_1$. Show that

- (i) $[T_1, T_2] = -[T_2, T_1]$.
- (ii) $[[T_1, T_2], T_3] + [[T_2, T_3], T_1] + [[T_3, T_1], T_2] = 0$.

Condition (ii) is called the Jacobi identity. Note that it implies that the commutation relation is not an associative multiplication, as in general it implies

$$[[T_1, T_2], T_3] - [T_1, [T_2, T_3]] = [[T_1, T_2], T_3] + [[T_2, T_3], T_1] = -[[T_3, T_1], T_2]$$

which need not be zero.

A vector space L with an additional bilinear operation $L \times L \rightarrow L$ satisfying conditions (i) and (ii) is called a *Lie algebra*. The elements need not come originally as a commutator of elements of an associative algebra, as they did here. But from a Lie algebra we can define the *enveloping algebra*, which is associative.

4 [5 pts] [Note: I don't expect you to completely resolve the difficulties here, but I want you to show that you recognize what the issues are, and give indications of how to resolve them.]

In quantum mechanics, the states of a system are taken to be a Hilbert space. For a single particle in one dimension, the wave functions $\psi(x)$ have an inner product given by $\langle \phi | \psi \rangle = \int_{\mathbb{R}} \phi^*(x)\psi(x) dx$. The classical degrees of freedom such as position x and momentum p are replaced by linear operators \mathbf{x} and \mathbf{p} or combinations of these, and the reality of these variables classically become conditions that the operators are *hermitian*, which means equal to its *adjoint*. The adjoint of a linear operator \mathbf{A} is a linear operator \mathbf{A}^\dagger such that $\langle \phi | \mathbf{A}\psi \rangle = \langle \mathbf{A}^\dagger \phi | \psi \rangle$ for any wave functions ψ and ϕ .

The expected value of a classical quantity represented by \mathbf{A} in a state represented by a normalized wavefunction ψ is given by $\langle A \rangle = \langle \psi | \mathbf{A}\psi \rangle = \int_{\mathbb{R}} \psi^* \mathbf{A}\psi dx$. Its complex conjugate is $\langle A \rangle^* = \langle \mathbf{A}\psi | \psi \rangle = \langle \psi | \mathbf{A}^\dagger \psi \rangle$, so it is real if \mathbf{A} is hermitian. The momentum operator is $\mathbf{p} = \frac{\hbar}{i} \frac{\partial}{\partial x}$, and then the commutator $[\mathbf{x}, \mathbf{p}] := \mathbf{x}\mathbf{p} - \mathbf{p}\mathbf{x}$ acting of a wave function $|\psi\rangle$ gives

$$x \frac{\hbar}{i} \frac{\partial \psi}{\partial x} - \frac{\hbar}{i} \frac{\partial}{\partial x} x \psi = i\hbar \psi \quad \text{so} \quad [\mathbf{x}, \mathbf{p}] = i\hbar$$

and \mathbf{x} and \mathbf{p} are canonically conjugate variables.

Clearly \mathbf{x} is hermitian, but how is it that \mathbf{p} is? That is, what have we been sloppy about in specifying our space of wave functions that needs specifying to verify that $\mathbf{p} = \mathbf{p}^\dagger$?

Perhaps it will be illuminating to consider instead the azimuthal angle ϕ and the z component of angular momentum, $L_z = \frac{\hbar}{i} \frac{\partial}{\partial \phi}$ with $[L_z, \phi] = i\hbar$. These have the same commutation relation as \mathbf{p} and \mathbf{x} . According to the Heisenberg uncertainty principle¹, if $[\mathbf{A}, \mathbf{B}] = i\mathbf{C}$, then the uncertainty in A, given by $(\Delta A)^2 = \int \psi^* (A - \langle A \rangle)^2 \psi dx$, and the uncertainty in B are restricted by $\Delta A \Delta B \geq \frac{1}{2} |\langle C \rangle|$. So $\Delta L_z \Delta \phi \geq \hbar/2$, but we can have states of definite L_z , so $\Delta L_z = 0$, while $-\pi \leq \phi \leq \pi$, so $\Delta \phi \leq \pi$. How can that be?

¹Proof: We need to use the Schwarz inequality:

$$\|\phi\|^2 \|\psi\|^2 \geq |\langle \phi | \psi \rangle|^2.$$

As $(\Delta A)^2 = \|(A - \langle A \rangle)\psi\|^2$ the Schwarz inequality tells us

$$\begin{aligned} (\Delta A)^2 (\Delta B)^2 &\geq | \langle (A - \langle A \rangle)\psi | (B - \langle B \rangle)\psi \rangle | = \langle \psi | (A - \langle A \rangle)(B - \langle B \rangle)\psi \rangle \\ &= \frac{1}{2} \left\langle \psi \left| \left[(A - \langle A \rangle)(B - \langle B \rangle) + (B - \langle B \rangle)(A - \langle A \rangle) + [A, B] \right] \psi \right\rangle \right|. \end{aligned}$$

The first two terms are hermitian conjugates so give a real value, while the commutator is i times the hermitian operator C , so gives an imaginary, and the absolute value must be \geq the imaginary part.